

## Invited Editorial Comment

### Twins and Twinning

Judith G. Hall

Department of Pediatrics and Medical Genetics, University of British Columbia, B.C. Children's Hospital, Vancouver, Canada

Twins have fascinated humanity since the beginning of recorded history. This issue of the *American Journal of Medical Genetics* includes a series of papers on twins and twinning. It is a timely and important contribution because in the last few years there have been many new observations concerning twins.

The usefulness of twins in studying the genetic contribution to various traits was first suggested by Galton [1875]. He proposed that by comparing the concordance of a specific trait or disorder in monozygotic twins (MZ) with dizygotic twins (DZ) it would be possible to distinguish between environmental and heritable effects (nurture vs. nature). Galton's suggestion led to the development of a body of knowledge which has been particularly useful in studying complex disorders [Bouchard et al., 1990; Phillips, 1993].

Recent technological advances in molecular and developmental genetics have uncovered a number of non-traditional mechanisms associated with discordance in MZ twins, e.g., mosaicism [Saul et al., 1990; Machin, 1995], imprinting, and uniparental disomy [Haskins-Olney et al., 1988]. These observations raise concerns about the premises upon which studies comparing MZ and DZ twins have been based. The study of the mechanisms leading to discordance in MZ twins and the role such mechanisms play in the human twinning process itself should provide new insights into early normal human development.

Prior to biochemical and molecular techniques, determination of zygosity was imprecise. Physical similarity was used for centuries. When it became clear that physical resemblance was not always an accurate discriminator, blood types and placentation [Benirschke, 1961] were used. However, blood types and serum proteins are not expected to always be discordant in DZ twins and placental membranes have occasionally been shown to be unreliable in establishing monozygosity [Bieber et al., 1981; Langlois et al., 1994, personal com-

munication]. Furthermore, a single placenta with vascular connections between the twins is expected in at least 70% of liveborn MZ twins [van Dijk et al., 1994; Machin et al., 1995]. Thus, blood type, serum proteins, and even DNA studies from either twin will be expected to show admixture or chimerism even if concordance does exist.

The Weinberg method is used to give a rough mathematical estimation of the proportion of MZ and DZ twins in the population [Weinberg, 1902]. It is based on the relative number of like-sex and unlike-sex twin births assuming that the excess of like-sex twins are MZ twins. Using this method the overall birth incidence of twins (both MZ and DZ) is estimated to be 1 in 80 births or 1 in 40 individuals in North America. The Weinberg method may be useful for statistical purposes; however, for accurate twin research, zygosity must be established using placental membranes and/or DNA studies which are ideally done in tissues other than blood [Hill and Jeffreys, 1985; Jeffreys et al., 1985; Akane et al., 1991].

The number of DZ twins estimated by the Weinberg method varies among different ethnic groups (high among blacks, low among the Japanese) [Imaizumi, 1990] and is considered to be related to elevated gonadotropin levels leading to multiple ovulation [Nylander, 1981]. In the last decade the incidence of liveborn DZ twins has increased in association with the use of fertility drugs and in vitro fertilization techniques [Derom et al., 1991]. By contrast, the incidence of MZ twins had been thought to be constant worldwide at 1 MZ twin birth every 330 births [Nylander, 1975]; however, the incidence of MZ twins has been reported to increase with artificially induced ovulation and in vitro fertilization techniques [Edwards et al., 1986; Derom et al., 1987].

At a recent international conference on twins, data from several countries suggest that the incidence of all twin births is increasing [Bressers et al., 1987; Bryan, 1994] independent of fertility drugs. Czeizel et al. [1994] examined the effect of multivitamins in preventing congenital anomalies; he reported that vitamin supplementation prior to and in early pregnancy increased the occurrence of twin livebirths (both MZ and DZ). This observation suggests that the effect of vitamins in the prevention of congenital anomalies may maintain

Received for publication October 11, 1994; revision received February 27, 1995.

Address reprint requests to Judith G. Hall, Department of Pediatrics and Medical Genetics, University of British Columbia, B.C. Children's Hospital, 4480 Oak St., Rm. 2D15, Vancouver, B.C., Canada V6H 3V4.

twin pregnancies, some of which would otherwise have been lost due to lethal congenital anomalies. Thus, the observed increase of liveborn twins in the general population may be related to better nutrition. The study of Czeizel et al. [1994] also shows an increase in recognizable miscarriages in the vitamin supplemented group, suggesting that even abnormal pregnancies may progress further into the pregnancy when the mother has nutritional supplementation.

With the development and use of ultrasonography in early human pregnancies, it has become clear that many more pregnancies start as twins than come to term delivery as twins. The term "vanishing twin" has been coined for those pregnancies which convert to singletons or are lost [Levi, 1976; Jeanty et al., 1981; Jauniaux et al., 1986]. Jauniaux et al. [1986] estimated that at least 70% of twin conceptions do not come to term as twins. A more recent report by Boklage [1990] estimated that multiple conceptions may constitute 12% of all pregnancies but that only 2% survive to term as twins. The incidence of twins among miscarriages is also high [Livingston and Poland, 1980]. The mechanism(s) by which one (or both) of the twins in a pregnancy abort or vanishes is not known. Some suggestions include postconceptional nondisjunction leading to chromosomal imbalance such as tetraploidy [Rudnicki et al., 1991], embryonic growth disorganization due to abnormal chromosomes [Livingstone and Poland, 1980], non-viable malformations in one or both twins, and disruptions such as vascular compromise of one or both twins [Landy et al., 1986]. It is not clear whether the "vanishing twin" phenomenon occurs primarily in MZ or DZ or equally in both types of twins.

An increase in most congenital anomalies is seen in one or both individuals of a twin pair [Källén, 1986]. In at least 10% of MZ twin pregnancies coming to birth, one or both twins has some type of congenital anomaly [Doyle et al., 1990]. These anomalies include malformations, disruptions, and deformations [Schinzel et al., 1979]. The true incidence of anomalies in MZ and DZ twins is hard to assess since many previous studies failed to determine zygosity correctly and some even assumed dizygosity if there was discordance because of the anomalies.

The cause of MZ twinning is not known. Familial MZ twinning (in which several relatives in a family have MZ twin pregnancies) is rare in humans [Harvey et al., 1977; Parisi et al., 1983; Shapiro et al., 1987a]. The cause of familial MZ twinning is thought to be related to a single gene possibly associated with inherited abnormalities of the zona pellucida or cell-to-cell connections which would allow cells to separate before implantation and placentation [Shapiro et al., 1987b]. Interestingly, no increase in congenital anomalies is apparent in familial MZ twinning. Abnormalities in the zona pellucida allowing early separation of cells, discordant X-inactivation in some female MZ twins [Burn et al., 1986; Goodship et al., 1984], or abnormalities in developmental clocks (i.e., time of fertilization, implantation, hatching, X-inactivation) [Boklage, 1987] have all been suggested for sporadic (non-familial) MZ twinning in humans.

Discordance has been observed frequently among MZ twins with disorders known to be genomically imprinted [Hall and Lopez-Rangel, 1995]. Discordance among MZ twins is also seen with mosaicism for chromosomal abnormalities, in many single gene (autosomal dominant and X-linked) disorders and with cytoplasmic inheritance. Based on these observations we have postulated that the development of a discordant cell line (by any mechanism including chromosomal, single gene mutation, mitochondrial mutation, uniparental disomy, somatic crossing over, X-inactivation, imprinting, etc.) early in development may be a cause for human MZ twinning (often without obvious physical discordance). In order to demonstrate discordance between MZ twins, it is important to remember that 70% of MZ twins who come to term have shared their circulation. Thus, examining white blood cell DNA may not demonstrate cytogenetic, cytoplasmic, and/or molecular discordance even though it exists. Fibroblasts (skin biopsy) is a much better tissue to look for discordance when studying MZ twins.

The study of twins and the twinning process will greatly benefit from the use of the new genetic technologies. Proper use of DNA techniques can provide unequivocal differentiation between MZ and DZ twins as well as uncovering a number of different types of genetic discordance. The availability and accuracy of these new techniques will allow new studies (or the repetition of old studies) involving twins to be conducted with the advantage of properly established zygosity. Hopefully subgrouping of twins from now on will not only be based on accurately diagnosed zygosity, but also on a description of placentation, placental connections, and molecularly defined discordance. There is much work to be done and much to be learned!

## REFERENCES

- Akane A, Matsubara K, Shiono H, Yamada M, Nakagome Y (1991): Diagnosis of twin zygosity by hypervariable RFLP markers. *Am J Med Genet* 41:96-98.
- Benirschke K (1961): Accurate recording of twin placentation. A plea to the obstetrician. *Obstet Gynecol* 18:334-347.
- Bieber FR, Nance MM, Morton CC, Brown JA, Redwine FO, Jordan RL, Mohanakumar T (1981): Genetic studies of an acardiac monster: Evidence of polar body twinning in man. *Science* 213:775-777.
- Boklage CE (1987): Invited editorial essay: Twinning, non-righthandedness and fusion malformations: evidence for heritable causal elements held in common. *Am J Med Genet* 28:67-84.
- Boklage CE (1990): Survival probability of human conceptions from fertilization to term. *Int J Fertil* 35:75-94.
- Bouchard TJ, Lykken DT, McGue M, Segal NL, Tellegen A (1990): Sources of human psychological differences, the Minnesota study of twins reared apart. *Science* 250:223-228.
- Bressers WMA, Eriksson AW, Kostense PJ (1987): Increasing trend in monozygotic twinning rate. *Acta Genet Med Gemellol* 36:397-399.
- Bryan E (1994): Trends in twinning rates. *Lancet* 343:1151-1152.
- Burn J, Povey S, Boyd Y, Munro EA, West L, Harper K, Thomas D (1986): Duchenne muscular dystrophy in one of monozygotic twin girls. *J Med Genet* 23:494-500.
- Czeizel A, Metneki J, Dudás I (1994): Higher rate of multiple births after periconceptual vitamin supplementation. *Lancet* 330:23-24.
- Derom C, Derom R, Vlietinck R, Van Den Berghe H, Thiery M (1987): Increased monozygotic twinning rate after ovulation induction. *Lancet* 1:1236.

- Derom R, Derom C, Vlietinck R, Van Maele G, Van Den Berghe H (1991): Twin pregnancies after medically assisted reproduction: epidemiology. *J Perinat Med* 19(Suppl):229-233.
- Doyle PE, Beral V, Botting B, Wale CJ (1990): Congenital malformations in twins in England and Wales. *J Epidemiol Community Health* 45:43-48.
- Edwards RG, Mettler L, Walters DE (1986): Identical twins and in vitro fertilization. *J In Vitro Fertil Embryo Trans* 3:114-117.
- Galton F (1875): The history of twins, as a criterion of the relative powers of nature and nurture. *J Anthropol Inst.* 12:566-576.
- Goodship J, Carter J, Speer A, Burn J (1994): X-inactivation patterns in monozygotic and dizygotic female twins. *Am J Med Genet* 61: 205-208.
- Harvey MAS, Huntley RMC, Smith DW (1977): Familial monozygotic twinning. *J Pediatr* 90:246-250.
- Haskins-Olney A, Buehler BA, Waziri L (1988): Wiedemann-Beckwith syndrome in apparently discordant monozygotic twins. *Am J Med Genet* 29:491-499.
- Hill AVS, Jeffreys AJ (1985): Use of minisatellite DNA probes for determination of twin zygosity at birth. *Lancet* 2:1394.
- Hall JG, Lopez-Rangel E (1995): Twins and twinning. In: AE Emery, DL Rimoin (eds): "Principles and Practice of Medical Genetics," 3rd ed. New York: Churchill Livingstone (in press).
- Imaizumi Y (1990): Triplets and higher order multiple births in Japan. *Acta Genet Med Gemellol* 39:295-306.
- Jauniaux E, Elkazen N, Leroy F (1986): Clinical and morphologic aspects of the vanishing twin phenomenon. *Obstet Gynecol* 68:577-581.
- Jeanty P, Rodesch F, Verhoogen C (1981): The vanishing twin. *Ultrasonics* 2:25-30.
- Jeffreys AJ, Wilson V, Thein SL (1985): Individual specific "fingerprints" of human DNA. *Nature* 316:76-77.
- Källén B (1986): Congenital malformations in twins: a population study. *Acta Genet Med Gemellol* 33:61-69.
- Landy HJ, Wiener S, Corson SL (1986): The "vanishing twin" ultrasonographic assessment of fetal disappearance in the first trimester. *Am J Obstet Gynecol* 155:14-20.
- Levi S (1976): Ultrasonic assessment of the high rate of human multiple pregnancy in the first trimester. *J Clin Ultrasound* 4:3-7.
- Livingston JE, Poland BJ (1980): A study of spontaneously aborted twins. *Teratology* 21:139-148.
- Machin G, Still K, Tasneem L (1995): Correlations of placental vascular anatomy and clinical outcomes in sixty-nine monochorionic twin pregnancies. *Am J Med Genet* 61:229-236.
- Machin GA (1995): Some causes of genotypic and phenotypic discordance in monozygotic twin pairs. *Am J Med Genet* 61:216-228.
- Nylander PPS (1975): Frequency of multiple births. In MacGillivray I, Nylander PPS, Corney G (eds): "Human Multiple Reproduction." London: WB Saunders, p 87.
- Nylander P (1981): The factors that influence twinning rates. *Acta Genet Med Gemellol* 30:189-202.
- Parisi P, Gatti M, Prinzi G, Caperna G (1983): Familial incidence of twinning. *Nature* 304:626-628.
- Phillips DW (1993): Twin studies in medical research. Can they tell us whether diseases are genetically determined? *Lancet* 341: 1008-1009.
- Rudniski M, Vejerslev LO, Junge J (1991): The vanishing twin: morphologic and cytogenetic evaluation of an ultrasonographic phenomenon. *Gynecol Obstet Invest* 31:141-145.
- Saul RA, Schwartz CE, Stevenson RE (1990): DNA evidence for somatic mosaicism in monozygotic twins discordant for Proteus syndrome. *Proc Greenwood Genet Center* 9:12-15.
- Schinzel AGL, Smith DW, Miller JR (1979): Monozygotic twinning and structural defects. *J Pediatr* 95:921-930.
- Shapiro LR, Zemek L, Shulman MJ (1978a): Genetic etiology for monozygotic twinning. New York: Alan R. Liss, Inc., for the National Foundation—March of Dimes BDOAS XIV(6A):219-225.
- Shapiro LR, Zemek L, Shulman MJ (1978b): Familial monozygotic twinning; An autosomal dominant form of monozygotic twinning with variable penetrance. In Nance W, Allen G, Parisi P (eds): "Twin Research: Biology and Epidemiology." New York: Alan R. Liss, pp 61-98.
- van Dijk BA, Boomsma DI, de Man AJM (1994): Blood group chimerism in human multiples is not rare. *Am J Med Genet* 61:264-268.
- Weinberg W (1902): Beiträge zur Physiologie und der Pathologie der Mehrlingsgeburten beim Menschen. *Arch Physiol* 88:346-430.